Surname	Other n	ames
Pearson Edexcel GCE	Centre Number	Candidate Number
Mechani	cs M1	
Advanced/Advan	ced Subsidiary	
Advanced/Advan Wednesday 8 June 2016 Time: 1 hour 30 minut	6 – Morning	Paper Reference 6677/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

P 4 6 7 0 8 A 0 1 2 8

Turn over ▶



1. [In this question i and j are horizontal unit vectors due east and due north respectively and position vectors are given relative to a fixed origin O.]

Two cars P and Q are moving on straight horizontal roads with constant velocities. The velocity of P is $(15\mathbf{i} + 20\mathbf{j})$ m s⁻¹ and the velocity of Q is $(20\mathbf{i} - 5\mathbf{j})$ m s⁻¹

(a) Find the direction of motion of Q, giving your answer as a bearing to the nearest degree.

(3)

At time t = 0, the position vector of P is 400**i** metres and the position vector of Q is 800**j** metres. At time t seconds, the position vectors of P and Q are **p** metres and **q** metres respectively.

- (b) Find an expression for
 - (i) \mathbf{p} in terms of t,
 - (ii) \mathbf{q} in terms of t.

(3)

(c) Find the position vector of Q when Q is due west of P.

-	4.	
1	4 N	
	+,	
•	-,	

uestion 1 continued	



Leave blank

Question 1 continued	

Question 1 continued	Leave
	Q1
(Total 10 marks)	
(-



2.




Figure 1

A vertical rope AB has its end B attached to the top of a scale pan. The scale pan has mass 0.5 kg and carries a brick of mass 1.5 kg, as shown in Figure 1. The scale pan is raised vertically upwards with constant acceleration 0.5 m s⁻² using the rope AB. The rope is modelled as a light inextensible string.

/ \	T 1	. 1	. •		. 1		4 D
(0)	Lind	tha	tension	110	tha	rana	$\Lambda \nu$
lai	THIU	uic	rension	111	uic	1000	AD
()						P -	

(3)

(b) I had the magnitude of the force exerted on the scare pair by the oriek.	(b)	Find the magnitude of the force exerted on the scale pan by the brick.	
--	-----	--	--

	Question 2 continued	blank
- $Q2$		\mathbb{Q}^2
(Total 6 marks)	(Total 6 marks)	



3. A particle P of mass 0.4 kg is moving on rough horizontal ground when it hits a fixed vertical plane wall. Immediately before hitting the wall, P is moving with speed 4 m s⁻¹ in a direction perpendicular to the wall. The particle rebounds from the wall and comes to rest at a distance of 5 m from the wall. The coefficient of friction between P and the ground is $\frac{1}{8}$.

Find the magnitude of the impulse exerted on *P* by the wall.

(7)



Question 3 continued	blank
Question 5 continued	
	Q3
(Total 7 marks)	



4. Two trains M and N are moving in the same direction along parallel straight horizontal tracks. At time t = 0, M overtakes N whilst they are travelling with speeds $40 \,\mathrm{m \, s^{-1}}$ and $30 \,\mathrm{m \, s^{-1}}$ respectively. Train M overtakes train N as they pass a point X at the side of the tracks.

After overtaking N, train M maintains its speed of $40 \,\mathrm{m\,s^{-1}}$ for T seconds and then decelerates uniformly, coming to rest next to a point Y at the side of the tracks.

After being overtaken, train N maintains its speed of 30 m s⁻¹ for 25 s and then decelerates uniformly, also coming to rest next to the point Y.

The times taken by the trains to travel between X and Y are the same.

(a) Sketch, on the same diagram, the speed-time graphs for the motions of the two trains between *X* and *Y*.

(4)

Given that XY = 975 m,

(b) find the value of *T*.

(8)

	Leave blank
Question 4 continued	



Leave blank

Question 4 continued	

	Leave blank
Question 4 continued	
	Q4
(Total 12 mai	rks)



5.




Figure 2

A particle P of mass 2 kg is held at rest in equilibrium on a rough plane by a constant force of magnitude 40 N. The direction of the force is inclined to the plane at an angle of 30° . The plane is inclined to the horizontal at an angle of 20° , as shown in Figure 2. The line of action of the force lies in the vertical plane containing P and a line of greatest slope of the plane. The coefficient of friction between P and the plane is μ .

Given that P is on the point of sliding up the plane, find the value of μ .	(

estion 5 continued	
	_



	Leave blank
Question 5 continued	

Question 5 continued	Le: bla
question 5 continued	



6. A non-uniform plank AB has length 6 m and mass 30 kg. The plank rests in equilibrium in a horizontal position on supports at the points S and T of the plank where AS = 0.5 m and TB = 2 m.

When a block of mass M kg is placed on the plank at A, the plank remains horizontal and in equilibrium and the plank is on the point of tilting about S.

When the block is moved to B, the plank remains horizontal and in equilibrium and the plank is on the point of tilting about T.

The distance of the centre of mass of the plank from A is d metres. The block is modelled as a particle and the plank is modelled as a non-uniform rod. Find

(i) the value of d,

(ii)	the	value	αf	M

(7)

	I
uestion 6 continued	
uestion o continueu	



Leave blank

Question 6 continued	
	_
	_
	_
	_
	-
	_
	-
	_
	_
	_
	_
	_
	_
	-
	_
	-
	_
	_
	_
	_
	_
	_
	_
	_
	-
	_

Question 6 continued		blank
		Q6
	(Total 7 marks)	
	(Iotai / marks)	



7. Two forces \mathbf{F}_1 and \mathbf{F}_2 act on a particle P.

The force \mathbf{F}_1 is given by $\mathbf{F}_1 = (-\mathbf{i} + 2\mathbf{j})$ N and \mathbf{F}_2 acts in the direction of the vector $(\mathbf{i} + \mathbf{j})$.

Given that the resultant of \mathbf{F}_1 and \mathbf{F}_2 acts in the direction of the vector $(\mathbf{i} + 3\mathbf{j})$,

(a) find \mathbf{F}_2

(7)

The acceleration of P is $(3\mathbf{i} + 9\mathbf{j})$ m s⁻². At time t = 0, the velocity of P is $(3\mathbf{i} - 22\mathbf{j})$ m s⁻¹

(b) Find the speed of P when t = 3 seconds.

(4)

uestion 7 continued	



Leave blank

Question 7 continued		

Question 7 continued		Leave blank
	Question 7 continued	
(Total 11 marks)		Q7
	(Total 11 marks)	



8.




Figure 3

Two particles P and Q have masses 1.5 kg and 3 kg respectively. The particles are attached to the ends of a light inextensible string. Particle P is held at rest on a fixed rough horizontal table. The coefficient of friction between P and the table is $\frac{1}{5}$. The string is parallel to the table and passes over a small smooth light pulley which is fixed at the edge of the table. Particle Q hangs freely at rest vertically below the pulley, as shown in Figure 3. Particle P is released from rest with the string taut and slides along the table.

Assuming that P has not reached the pulley, find

(a) the tension in the string during the motion,

(8)

(b) the magnitude and direction of the resultant force exerted on the pulley by the string.



	Leave
	blank
Question 8 continued	

